

# Example: 3 mL of Diluted CCH Remaining After Treating the Buttocks\*

1. Dilution: You need **3 mL x 4** for a total of 12 mLs of sterile, non-preserved saline.


Find CCH Remaining  
On the **Dilution Table**

Amount of Reconstituted CCH x 4	mL of Sterile Saline to Add	Total mL for Injection
1 mL x 4 =	4 mL	5 mL
2 mL x 4 =	8 mL	10 mL
<b>3 mL x 4 =</b>	<b>12 mL</b>	<b>15 mL</b>
4 mL x 4 =	16 mL	20 mL
5 mL x 4 =	20 mL	25 mL
6 mL x 4 =	24 mL	30 mL

2. Using a 20 mL vial of sterile, non-preserved saline, **remove 8 mL and discard.**

Remove  
**8 mL**  
Saline

Leave  
**12 mL**  
Saline



20mL Vial Sterile,  
Non-Preserved Saline


3. To the remaining 12 mL of sterile, non-preserved saline, **add the remaining 3 mL of reconstituted CCH.**

Add **3 mL**  
CCH Back to  
the Saline



4. This gives you **15mL of diluted CCH**, which is enough to inject 10 thigh dimples, using the 5-point technique.

This Yields  
**15 mL**  
Diluted CCH



\*Always use sterile, non-preserved saline for dilution of CCH